

DEEP POCKETS IN LATTICES AND OTHER GROUPS

ANDREW D. WARSHALL

ABSTRACT. We show the nonexistence of deep pockets in a large class of groups, extending a result of Bogopol'skii. We then give examples of important groups (namely lattices in Nil and Sol) which have deep pockets.

1. INTRODUCTION

Let G be an infinite group and A a finite generating set for G . We define the *depth* (or more verbosely the *dead-end depth*) of an element $g \in G$ with respect to A to be the distance (in the word metric with respect to A) from g to the complement of the radius- $d_A(1, g)$ closed ball about the identity in G . In fact this definition makes sense if G is replaced with an arbitrary pointed metric space; this will be used in Section 3. Depth must be finite for all $g \in G$ since G is infinite and finitely generated. If the depth of g is > 1 then g is called a *dead end*. (The idea is that a geodesic word representing g cannot be extended further if it is to remain a geodesic.) We define the depth of G with respect to A to be the supremum of the depths of all elements of G .

A natural question to ask is whether the depth of G for some given generating set A is finite; if it is not, G is said to have *deep pockets* with respect to A . One result in this direction has been proven by Bogopol'skii (see [1]), who showed that all hyperbolic groups have finite depth with respect to all generating sets. We extend this result to a broader class of groups, namely those with a regular language of geodesics. To be precise, we prove the following

Proposition 1. *Let G be an infinite group with a regular language L of geodesics with respect to A a finite generating set for G . Then there exists a uniform bound for the depth of elements of G with respect to A .*

Definition 1. A group is said to be *weakly geodesically automatic* if it admits an automatic structure made up entirely of geodesics.

Corollary 2. *Any group G weakly geodesically automatic with respect to a generating set A has a uniform bound on depth with respect to A .*

Two cases of this corollary are hyperbolic groups and finitely generated abelian groups, each with respect to any set of generators; for details see [4]. The first of these coincides with Bogopol'skii's result.

The next proposition provides a slight extension of the finitely generated abelian case.

Date: February 2, 2008.

2000 Mathematics Subject Classification. 20F65.

We thank our advisor, Andrew Casson, John Mackay and Joshua Zelinsky for their helpful comments.

Proposition 3. *Any euclidean group has a uniform bound on depth with respect to any set of generators.*

These propositions seem unsurprising, since previous constructions of groups with deep pockets tended to involve wreath products or the like; see [3], [2] and [5]. However, we here give two examples of comparatively well-known and well-behaved groups whose depth is infinite with respect to reasonably standard generating sets.

Proposition 4. *Let $H = \langle a, b \mid [a, [a, b]], [b, [a, b]] \rangle$ be the discrete Heisenberg group. Then H has infinite depth with respect to its two-element generating set $\{a, b\}$.*

Proposition 5. *Let R be a hyperbolic automorphism of $\mathbb{Z}^2$ and let $G_R = \mathbb{Z}^2 \rtimes_R \mathbb{Z}$. Let $\{a, b\}$ be the standard generating set for $\mathbb{Z}^2$ and $\{c\}$ be that for $\mathbb{Z}$. Then G_R has arbitrarily deep dead ends with respect to $\{a, b, c\}$.*

It is interesting that H and G_R are lattices in two of Thurston's eight three-dimensional model geometries (see [6]), namely Nil and Sol respectively. In contrast, it follows from Proposition 1 (or [1]) and Proposition 3 that no lattice in $\mathbb{H}^3$, $S^2 \times \mathbb{R}$, $\widehat{SL_2(\mathbb{R})}$ or $\mathbb{E}^3$ can have deep pockets. Since the question of admitting a lattice with deep pockets is meaningless for S^3 since it is compact (so that any lattice must be finite), there remains one model geometry for which it is (to the best of this author's knowledge) open, namely $\mathbb{H}^2 \times \mathbb{R}$.

2. PROOF OF PROPOSITION 1

Proof. Since G is infinite, every individual element has finite depth. Let n be the number of states of a (deterministic) finite state automaton F accepting L . Then it will suffice to prove the existence of a uniform bound on depth for elements at distance at least n from the identity in G with respect to A . Let g be such an element and consider the sequence of states assumed by F as it reads the element w of A representing g . This sequence has length at least $n+1$, so, by the pigeonhole principle, the last $n+1$ terms of this sequence must contain at least one repetition. We thus express w as abc , where $|bc| \leq n$, $|b| > 0$ and F is in the same state after reading ab as after reading a . Then F will also accept $abbc$, since after reading abb it will be in the same state as after reading ab , so that reading c will get it into an accept state. Since all words accepted by F are geodesic, $abbc$ is thus a geodesic word of length greater than that of $w = abc$ (since $|b| > 0$) and representing a group element within $|c| + |bc| \leq 2|bc| \leq 2n$ of g . Thus the depth of g is $\leq 2n$, so we are done. $\square$

3. PROOF OF PROPOSITION 3

We need the following auxiliary result, which will also be used in Section 5.

Proposition 6. *Let f be a function from a metric space A to $\mathbb{Z}$ and $n \in \mathbb{Z}$. Suppose there exists $a \in A$ and $r \in \mathbb{Z}$ such that for all $a' \in B_r(a)$ $f(a') \leq f(a) + n$. Then there exists some $a' \in A$ such that f attains a maximum on $B_{r/n}(a')$ at a' .*

Proof. Define $g(x) = \max_{b \in B_a(x)} f(b)$ for $x > 0$, $f(a)$ for $x = 0$. Clearly, g is nondecreasing and, by hypothesis, $g(r) \leq f(a) + n = g(0) + n$. Thus there must exist some x such that g is constant on the interval $(x, x + r/n]$, where $x + r/n \leq r$. Then we may just set a' to be an element of A at distance $\leq x$ from a such that $f(a') = g(x)$. By the triangle inequality and the definition of g , f is bounded on $B_{r/n}(a')$ by $g(x + r/n) = g(x) = f(a')$. $\square$

From this we deduce

Proposition 7. *Suppose a pointed metric space (A, a_0, d) has deep pockets. Then if C is a real number and d' is a integer-valued metric on A such that*

$$|d'(a_1, a_2) - d(a_1, a_2)| < C$$

for all $a_1, a_2 \in A$ then the resulting pointed metric space (A, a_0, d') will still have deep pockets.

Proof. Let $f: A \rightarrow \mathbb{Z}$ be distance from the origin with respect to the new metric, and a have depth at least $r + C$ with respect to (A, a_0, d) . Then Proposition 6 gives a dead end with respect to (A, a_0, d') of depth at least r/C . Since r was arbitrary, we are done. $\square$

With these preliminaries out of the way, we approach the heart of the proof, starting with the following

Definition 2. We define a *weighted* generating set for $\mathbb{Z}^n$ to be an ordered pair (A, μ) , where A is a generating set for $\mathbb{Z}^n$ and $\mu: A \rightarrow \mathbb{N}$ is called the *weight* function. We say that the *length* of the word $a_1 a_2 \dots a_m$ with respect to the weight function μ is $\sum_{i=1}^m \mu(a_i)$.

It is clear that a weighted generating set defines a word metric just as does an ordinary generating set.

Proposition 8. *Every finite weighted generating set for $\mathbb{Z}^n$ has depth bounded for all group elements.*

Let (A, μ) be a generating set for $\mathbb{Z}^n$. Let M be the least common multiple of the elements of the image of μ and let A' be the set consisting of $aM/\mu(a)$ for every $a \in A$. If we regard $\mathbb{Z}^n$ as being embedded in the obvious way as a lattice in $\mathbb{R}^n$ and let B be the convex hull of $A' \cup A'^{-1}$ in $\mathbb{R}^n$, then B will clearly be a convex polytope. In fact, its dimension will be n , since otherwise A would be contained in an $n - 1$ -dimensional subspace, hence could not generate. We have the following

Proposition 9. *If $a_1, a_2, \dots, a_m \in A' \cup A'^{-1}$ are the vertices of a facet of B and $i_1, i_2, \dots, i_m$ are nonnegative integers then $w = a_1^{i_1} a_2^{i_2} \dots a_m^{i_m}$ is a geodesic word. In fact, w lies on a geodesic ray.*

Proof. The facet of B of which $a_1, a_2, \dots, a_m$ are vertices must lie in some hyperplane H in $\mathbb{R}^n$, say $\mathbf{a} \cdot \mathbf{x} = b$. Since B is symmetric about the origin and lies in one of the closed half-spaces bounded by H , if the origin lay in H then B would lie entirely in H , so so would $A' \cup A'^{-1}$, so $A \cup A^{-1}$ would as well, which is a contradiction since A generates $\mathbb{Z}^n$. Thus $b \neq 0$, so we may choose $\mathbf{a}$ so that $b = 1$, so H is given by the equation $\mathbf{a} \cdot \mathbf{x} = 1$ and B lies in the closed half-space given by $\mathbf{a} \cdot \mathbf{x} \leq 1$. In particular, so does $A' \cup A'^{-1}$, so if $g \in \mathbb{Z}^n$ is given by a word of length l with respect to (A, μ) then $\mathbf{a} \cdot g \leq l/M$. Setting $g = a_1^{i_1} a_2^{i_2} \dots a_m^{i_m}$, we have $\mathbf{a} \cdot g = \sum_{j=1}^m i_j$, so the minimal length for a word representing g is $M \sum_{j=1}^m i_j$. But this is the length of w since each a_i has length M , proving the first sentence. The second sentence follows clearly. $\square$

Proposition 10. *Every element of $\mathbb{Z}^n$ is within bounded distance (with respect to (A, μ)) of a geodesic ray.*

Proof. By choosing a finite triangulation of each facet f_i of B , we can assume WLOG that they are all $n - 1$ -simplices, hence have n vertices. As in the proof of Proposition 9, no facet may be confined to any hyperplane through the origin, so the vertices of each facet (since they clearly cannot lie in one $n - 2$ -plane) must be linearly independent, hence generate a maximal-rank lattice. If we regard this lattice as a based lattice embedded in $\mathbb{R}^n$ with the usual norm, it will partition $\mathbb{R}^n$, hence $\mathbb{Z}^n$, into closed parallelepipeds. Clearly there exists some bound d_i on the A -distance between any element of $\mathbb{Z}^n$ contained in the closed unit parallelepiped generated by the facet f_i and the origin. By translation, we find that every element of $\mathbb{Z}^n$ is within d_i of some corner of every parallelepiped in which it lies. Since B has finitely many facets, we may set d to be the greatest of the d_i .

I claim that every element of $\mathbb{Z}^n - B$ is within d of a geodesic ray. So let $g \in \mathbb{Z}^n - B$, and consider the line segment (in $\mathbb{R}^n$) s between g and the origin. Since the origin is in B , $g \notin B$ and B is convex, s intersects the boundary of B exactly once (say at t), so we let $a_1, a_2, \dots, a_n$ be the vertices of a (closed) facet containing this intersection. By the last paragraph, g is within distance d of some corner of every parallelepiped of the $a_1 a_2 \dots a_n$ -lattice to which it belongs. But the coordinates of g with respect to $a_1, a_2, \dots, a_n$ as an element of $\mathbb{R}^n$ must be nonnegative since they are obtained by multiplying a positive number by the coordinates of t , which is (as a vector) a weighted average of $a_1, a_2, \dots, a_n$. Hence g belongs to a parallelepiped all of whose corners have all their coordinates nonnegative, hence (by Proposition 9) lie on a geodesic ray, proving our claim.

Since B is bounded, the case $g \in B$ is trivial. $\square$

Proof of Proposition 8. Let D be the bounded distance whose existence was proven in the preceding proposition. Let $g \in \mathbb{Z}^n$ with minimal word length l . Then g is at distance at most D from an element on a geodesic ray whose (A, μ) -distance from the origin is, by the triangle inequality, at least $l - D$. Then there exists a group element at distance at least $D + 1$ but at most $D + M + 1$ further out along this ray whose distance from the origin is at least $l + 1$ and, by the triangle inequality again, at distance at most $2D + M + 1$ from g . Thus g has depth at most $2D + M + 1$, as claimed. $\square$

Let A be a generating set on E a euclidean group and let $\mathbb{Z}^n \triangleleft E$. Let $\phi: F_A \twoheadrightarrow E$ be the projection map. We denote the length of a word $w \in F_A$ by $l(w)$. Let A' be the set of words w in $A \cup A^{-1}$ such that $\phi(w) \in \mathbb{Z}^n$ but, for any proper subword w' of w , $\phi(w') \notin \mathbb{Z}^n$. Let (B, μ) be the finite weighted generating set for $\mathbb{Z}^n$ where $B = \{a^g \mid a \in A'\} g \in E = \bigcup_{h \in E/\mathbb{Z}^n} A'^h$ and μ is defined so that $\mu(a^g)$ is the length of a (as a word in A) for $a \in A'$ and $g \in E$. It will follow from Proposition 11 that this is in fact a generating set. In any event, let $\pi: \mathbb{Z}^B \rightarrow \mathbb{Z}^n$ be the projection and let $l_\mu(v)$ denote the length of the word $v \in \mathbb{Z}^B$ with respect to μ . We denote the distance of $x \in \mathbb{Z}^n$ from the origin (at the moment, possibly infinite) with respect to this generating set by $\|x\|$.

Proposition 11. *Suppose $w \in F_A$ with $\phi(w) = x \in \mathbb{Z}^n$. Then $\|x\| \leq l(w)$. In particular, B is a generating set for $\mathbb{Z}^n$.*

Proof. Since $\phi(w) \in \mathbb{Z}^n$, w must contain some minimal subword under inclusion in $\phi^{-1}(\mathbb{Z}^n)$, say w_1 . By definition, $w_1 \in A'$. Then $w = w_l w_1 w_r$ may be replaced with

$w_1^{w_l^{-1}} w_l w_r$, where $w_1^{w_l^{-1}} \in B$ and

$$\mu(w_1^{w_l^{-1}}) = \mu(w_1) = l(w_1) = l(w) - l(w_l) - l(w_r) \leq l(w) - l(w_l w_r)$$

by definition and the triangle inequality. The first claim is now proven by induction. The second claim follows easily. $\square$

Proposition 12. *There exists a D such that for any $x \in \mathbb{Z}^n$ there is a word $w \in F_A$ with $\phi(w) = x$ and $l(w) \leq \|x\| + D$.*

Proof. Let $m = [E : \mathbb{Z}^n]$ and pick a set of words $w_1, \dots, w_m$ in F_A representing each of the finitely many cosets of $\mathbb{Z}^n$ in E . Consider a word $v \in \mathbb{Z}^B$ with $\pi(v) = x$ and $l_\mu(v) = \|x\|$. Since $\mathbb{Z}^n$ is abelian, we can assume that v is ordered to separate out the letters in each A^h . The word v is thus divided into m pieces $v_1 v_2 \dots v_m$, where each $v_i \in \mathbb{Z}^{A^h}$ for some specific $h \in E/\mathbb{Z}^n$. Thus each v_i can be replaced with $w_i^{-1} v'_i w_i$ for some $v'_i \in \mathbb{Z}^{A'}$ with $l_\mu(v'_i) = l_\mu(v_i)$. But then each v'_i can be replaced with a $v''_i \in F_A$ with $l(v''_i) \leq l_\mu(v'_i) = l_\mu(v_i)$ by construction of A' and μ . We thus have

$$l(w_i^{-1} v''_i w_i) \leq 2l(w_i) + l(v''_i) \leq 2l(w_i) + l_\mu(v_i).$$

Concatenating these yields $w = w_1^{-1} v''_1 w_1 w_2^{-1} v''_2 w_2 \dots w_m^{-1} v''_m w_m \in F_A$ with $\phi(w) = x$ and

$$\begin{aligned} l(w) &\leq \sum_{i=1}^m l(w_i^{-1} v''_i w_i) \leq \sum_{i=1}^m (2l(w_i) + l_\mu(v_i)) \\ &= 2 \sum_{i=1}^m l(w_i) + l_\mu(v) = 2 \sum_{i=1}^m l(w_i) + \|x\|, \end{aligned}$$

so we are done if we set $D = 2 \sum_{i=1}^m l(w_i)$. $\square$

Proof of Proposition 3. It follows from Propositions 8, 11, 12 and 7 that $\mathbb{Z}^n$ cannot have deep pockets with respect to the subspace metric induced from the metric with respect to A on E . But since $[E : \mathbb{Z}^n] < \infty$, every element of E lies within bounded distance b of $\mathbb{Z}^n$. Thus if E had a dead end of depth d then $\mathbb{Z}^n$ would have (with respect to the subspace metric) a dead end of depth at least $d/b + 1$, by Proposition 6. Thus E cannot have deep pockets, as claimed. $\square$

4. PROOF OF PROPOSITION 4

We now turn our attention to the Heisenberg group H with respect to its standard two-generator presentation $\langle a, b \mid [a, [a, b]], [b, [a, b]] \rangle$. (We use $[x, y]$ to denote $x^{-1}y^{-1}xy$.) It is well known that every element of H is expressible uniquely as a word of the form $a^i b^j [a, b]^k$, $a, b, c \in \mathbb{Z}$. In general, of course, these words are not of minimal length; for example, ba is expressed as $ab[a, b]^{-1}$. However, we give the following way of visualizing elements of H , from which insight can be gained as to minimal-length words.

Consider i and j as coordinates in the standard integer lattice $\mathbb{Z}^2$. Then we can represent a word in a and b as a path in $\mathbb{Z}^2$ starting at the origin. Multiplication on the right by $a^{\pm 1}$ corresponds to appending to the path a segment one unit to the right or left, while multiplication by $b^{\pm 1}$ corresponds to appending a segment one unit up or down. It is clear that this gives a one-to-one correspondence between

words and paths and that a word representing $a^i b^j [a, b]^k$ corresponds to a path from the origin to (i, j) . If to this path we append a vertical segment leading from (i, j) to $(i, 0)$ and a horizontal segment from $(i, 0)$ to the origin, we get a loop based at the origin. This loop will correspond to a word representing $a^i b^j [a, b]^k b^{-j} a^{-i} = a^i b^j b^{-j} a^{-i} [a, b]^k = [a, b]^k$. We have thus reduced the study of the correspondence between paths starting at the origin and elements of H to that of the correspondence between loops based at the origin and powers of $[a, b]$.

Proposition 13. *If two loops based at the origin correspond to the same power of $[a, b]$, then they enclose the same oriented area.*

Proof. Consider two loops based at the origin that correspond to the same power of $[a, b]$. Then one can be transformed into the other by appending loops conjugate to $a^{-1}a$, aa^{-1} , $b^{-1}b$, bb^{-1} , $[a, [a, b]]$, $[b, [a, b]]$, $[[a, b], a]$ and $[[a, b], b]$. But the area enclosed by the concatenation of two loops is the sum of the areas enclosed by each one, so it suffices to show that any loop conjugate to one of the eight listed above encloses zero area. Since conjugation does not affect the area enclosed by a loop, it in fact suffices to show that the eight loops listed above enclose zero area. But this can be checked trivially. $\square$

Proposition 14. *Any loop based at the origin corresponds to $[a, b]^k$, where k is the oriented area it encloses.*

Proof. By Proposition 13 and the discussion preceding it, it suffices to show that the path corresponding to the word $[a, b]$ encloses an oriented area of 1. But this is clear. $\square$

Proposition 15. *Any path P from the origin to (i, j) corresponds to $a^i b^j [a, b]^k$, where k is the oriented area enclosed by the concatenation of P and the path corresponding to $b^{-j} a^{-i}$.*

Proof. This is just the combination of Proposition 14 and the discussion preceding Proposition 13. $\square$

Proposition 16. *The element $[a, b]^{n^2+1}$ is at distance $4n + 2$ from the identity.*

Proof. Clearly, $[a, b]^{n^2+1}$ is within $4n + 2$ of the identity, since it equals

$$a^{-n-1} b^{-1} a b^{-n+1} a^n b^n.$$

So consider a word w representing $[a, b]^{n^2+1}$. By Proposition 15, we know that w corresponds to a loop based at the origin enclosing oriented area $n^2 + 1$. If we let x be the measure of the projection of the loop to the horizontal axis and y the measure of that to the vertical axis, we have $xy > n^2$, so, by the arithmetic mean-geometric mean inequality, $x + y > 2n$, so the length of w is $> 4n$. Since w corresponds to a loop, its length must be even, so in fact it is at least $4n + 2$. $\square$

Proposition 17. *The element $a^i b^j [a, b]^k$ with $|i| < n$, $|j| < n$, $|k| < n(n + 1)$ is within $4n + 2$ of the identity.*

Proof. Since, by Proposition 15, rotations of the plane by $\pi/2$ about the origin and reflections about the line $j = 0$ do not affect distance from the identity, we may assume WLOG that $i \geq |j|$ and $k \geq 0$. Let $k = q(n + 1) + r$, $0 \leq r \leq n$, $0 \leq q \leq n - 1$. Then clearly $b^{-q-1} a^r b a^{n+1-r} b^q a^{-n-1}$ is a loop based at the origin of length $2q + 2n + 4 \leq 4n + 2$ bounding oriented area $n(n + 1)$. Thus

$b^{-q-1}a^rba^{n+1-r}b^qa^{i-n-1}b^j$ represents $a^ib^j[a, b]^k$, and it will have length $\leq 4n+2$ since $|j| \leq i \leq n+1$. $\square$

Proposition 18. *Every element within distance m of $[a, b]^{n^2+1}$ is of form*

$$a^ib^j[a, b]^k$$

with $|i| \leq m$, $|j| \leq m$ and $|k| \leq n^2 + 1 + m(m-1)/2$.

Proof. We proceed by induction. The statement is clear for $m = 0$. So consider $w = a^ib^j[a, b]^k$ with $|i| \leq m$, $|j| \leq m$ and $|k| \leq n^2 + 1 + m(m-1)/2$. Then $wb = a^ib^{j+1}[a, b]^k$, $wb^{-1} = a^ib^{j-1}[a, b]^k$, $wa = a^{i+1}b^j[a, b]^{k-j}$ and $wa^{-1} = a^{i-1}b^j[a, b]^{k+j}$, so, by the triangle inequality, these four new words can all be expressed as $a^{i'}b^{j'}[a, b]^{k'}$ with $|i'| \leq m+1$, $|j'| \leq m+1$ and $|k'| \leq n^2 + 1 + m(m-1)/2 + |j| \leq n^2 + 1 + m(m-1)/2 + m = n^2 + 1 + m(m+1)/2$, as claimed. $\square$

Proposition 19. *If $n > 2$, the element $g_n = [a, b]^{n^2+1}$ is a dead end of depth at least $\sqrt{2n-4} + 1$. In particular, H has arbitrarily deep dead ends with respect to the generating set $\{a, b\}$.*

Proof. By Proposition 16, g_n is at distance $4n+2$ from the identity. But, by Propositions 17 and 18, any group element within $\sqrt{2n-4}$ of g_n is within $4n+2$ of the identity. The second sentence then follows clearly. $\square$

5. PROOF OF PROPOSITION 5

Let R be a hyperbolic automorphism of $\mathbb{Z}^2$. We turn our attention to the group

$$G_R = \mathbb{Z}^2 \rtimes_R \mathbb{Z}.$$

We denote the standard generators of $\mathbb{Z}^2$ by a and b and that of $\mathbb{Z}$ by c . This group is soluble, as its commutator is contained in the span of a and b , which commute with each other. Thus $G'_R = \langle a, b \rangle \cong \mathbb{Z}^2$ is a normal subgroup and $G_R/G'_R = \langle G'_R c \rangle$. The powers of c are thus a complete set of coset representatives, so every element of G_R can be expressed uniquely in the form $a^ib^jc^k$, where i, j and $k \in \mathbb{Z}$. We will use a, b and c to denote these three elements of G_R throughout.

Let F_S be the free group on the set S and let N be the normal closure of a and b in F_S . We know N is freely generated by $\{a^{c^i}, b^{c^i} \mid i \in \mathbb{Z}\}$. Any element $g \in F_S$ can be expressed uniquely as a product uc^i , where $u \in N$ and $i \in \mathbb{Z}$. Define $\phi: F_S \rightarrow N_{ab}$ to send g to the image of u under the abelianization map. (We denote by N_{ab} the abelianization of the group N .) Note that N_{ab} is freely generated as an abelian group by $\{a^{c^i}, b^{c^i} \mid i \in \mathbb{Z}\}$.

Define $\pi: N_{ab} \rightarrow G'_R$ to send $v \in N_{ab}$ to the element of G'_R represented by v . Similarly, define $\sigma_R: F_S \rightarrow G_R$ to send $w \in F_S$ to the element of G_R represented by w .

Notation. If $w \in F_S$ or N_{ab} , let $l(w)$ denote its length in the given sets of generators.

We can now state the following proposition, which is central to the proof of Proposition 5 and whose proof we postpone.

Proposition 20. *Let $G_R = \mathbb{Z}^2 \rtimes_R \langle c \rangle$, where R is a hyperbolic automorphism of $\mathbb{Z}^2$, and let $S = \{a, b, c\}$, where a and b are the standard generators of $\mathbb{Z}^2$. Then there exists D with the following property. Let $g = uc^n \in G_R$, where $u \in G'_R$ and*

$n \in \mathbb{Z}$. Let $v \in N_{ab}$ be a minimal-length element of $\pi^{-1}(u)$. Then there is some $w \in F_S$ with $\phi(w) = v$, $\sigma_R(w) = g$ and

$$l(w) - D \leq |g| \leq l(w),$$

where by $|g|$ we mean the length of the minimal-length element of $\sigma_R^{-1}(g)$.

Definition 3. Let $g = uc^z \in G_R$ with $u \in G'_R$. Let $\|g\|$ represent the minimal length of any element w of $\sigma_R^{-1}(g)$ such that $\phi(w)$ is a minimal-length element of $\pi^{-1}(u)$.

Remark. Any element $w \in \sigma_R^{-1}(g)$ must satisfy $\pi(\phi(w)) = u$.

In this notation, Proposition 20 implies that there exists a D dependent only on R such that, for any $g \in G_R$, $\|g\| - D \leq |g| \leq \|g\|$.

Notation. Let u_1 denote a and u_2 denote b .

The following lemma, an adaptation of Cleary and Taback's result on word length in wreath products, is used both in proving and in applying Proposition 20.

Lemma 21. Let $g = uc^z \in G_R = \mathbb{Z}^2 \rtimes_R \langle c \rangle$, where R is a hyperbolic automorphism of $\mathbb{Z}^2$ and $u \in G'_R$. Let $v = \sum_{j=1}^n u_{k_j}^{c^{i_j}} \in N_{ab}$ (where all the k_j are 1 or 2) such that $\pi(v) = u$ and let $S = \{a, b, c\}$ where a and b are the standard generators of $\mathbb{Z}^2$. Then the minimal length of all words $w \in F_S$ such that $\sigma_R(w) = g$, $\phi(w) = v$ is

$$2(\max(i_j, 0) - \min(i_j, 0)) + \min(|z - \max(i_j, 0)| - \max(i_j, 0), |z - \min(i_j, 0)| + \min(i_j, 0)) + n.$$

Proof. Consider the natural maps π_S and π_T mapping F_S and $\mathbb{Z}^T$, respectively, to

$$\mathbb{Z}^2 \wr \mathbb{Z} = \langle a, b, c \mid [a^{c^i}, b^{c^j}], [a^{c^i}, a^{c^j}], [b^{c^i}, b^{c^j}] \rangle.$$

A word $w \in F_S$ satisfies $\sigma_R w = g$ and $\phi w = v$ if and only if $\pi_S(w) = \pi_T(v)c^z$ in $\mathbb{Z}^2 \wr \mathbb{Z}$. (This is because $\phi(w) = v$ if and only if $\pi_S(w) = \pi_T(v)c^k$ for some k , and then we have $\sigma_R(w) = \pi(v)c^k$.) Thus the minimal length for w is simply the minimal length of a word in F_S representing $\pi_T(v)c^z$ in $\mathbb{Z}^2 \wr \mathbb{Z}$. The result now follows from Cleary and Taback's length formula in [3] for wreath products $G \wr \mathbb{Z}$, G some group, with respect to a generating set consisting of generators of G and a standard generator for $\mathbb{Z}$. $\square$

The final main ingredient is the following proposition, which tells us about the geometry of the $\|\cdot\|$ norm and whose proof we likewise postpone. Note that, for any group G with generating set A , d_A refers to the word metric on G induced by A .

Proposition 22. Let $G_R = \mathbb{Z}^2 \rtimes \mathbb{Z}$, where R is a hyperbolic automorphism of $\mathbb{Z}^2$, and let a and b be the standard generators of $\mathbb{Z}^2$. Let n be a positive integer. Then for all sufficiently large positive integers m there exists $B_{m,n} \subset G'_R$ such that every $u \in B_{m,n}$ has the following properties:

- If u is expressed as $a^i b^j$, $i, j \in \mathbb{Z}$, then $|i|, |j| < |\text{tr } R|^m$ and
- If $v \in N_{ab}$ is a minimal-length element of $\pi^{-1}(u)$ then $M(v) > n$ and $m(v) < -n$.

Furthermore, there exists $p = p(R, m, n) \in \mathbb{Z}$ and $u \in B_{m,n}$ such that for all u' with $d_{\{a,b\}}(u, u') \leq p$ the following conditions hold:

- $g' \in B_{m,n}$ and
- $\|u'\| \leq \|u\|$.

Finally, $\lim_{m \rightarrow \infty} p(R, m, n) = \infty$ for every R and n .

With these ingredients, we give the

Proof of Proposition 5. By Proposition 20, there exists D such that, for any $g \in G_R$, $\|g\| - D \leq |g| \leq \|g\|$. Thus to prove that G_R has deep pockets with respect to the word metric in S it suffices to prove that for any r we can find an element $h \in G_R$ such that $\| \cdot \|$ attains a maximum on $B_r(h)$ at h , for we may then apply Proposition 6 with $f(g) = |g|$ and $n = D$.

To that end, we choose $n \geq r$ and then m such that $r\tau^r \leq p(R, m, n)$, where p is defined as in Proposition 22; this choice of m is possible since $\lim_{m \rightarrow \infty} p(R, m, n) = \infty$. Define $u \in G'_R$ to be the element given by Proposition 22 with $B_{m,n}$ as input. Then let $g = u'c^i \in G_R$, $u' \in G_R$, be any element within distance r of u with respect to S . Then $|i| \leq r$. Furthermore, if $s \in G'_R$, $sc^{i'}a^{\pm 1} = sR^{-i'}(a^{\pm 1})c^{i'}$ and similarly if a is replaced by b , so right multiplication of any $u''c^{i'}$, $u'' \in G'_R$, $|i'| \leq r$, by any letter of S or its inverse changes u'' (with respect to $\{a, b\}$) by at most $|\tau|^r$. Thus $d_{\{a,b\}}(u, u') < r\tau^r \leq p(R, m, n)$, so $u' \in B_{m,n}$. By Proposition 22 and Lemma 21, this implies that $\|g\| \leq \|u'\|$. But Proposition 22 also tells us that this is $\|u'\| \leq \|u\|$, so we are done. $\square$

It remains to prove Propositions 20 and 22. Let p_R be the characteristic polynomial of R . We define $\Lambda_R = \mathbb{Z}[t]/(p_R(t))$. There is a natural map $\alpha_R: \mathbb{Z}[t, t^{-1}] \rightarrow \Lambda_R$, where by $\mathbb{Z}[t, t^{-1}]$ we mean the ring of Laurent polynomials in one variable t over $\mathbb{Z}$. Define $\tau = \alpha_R(t) \in \Lambda_R$. By abuse of notation, we also denote the greater (in absolute value) of the two eigenvalues of R by τ . Any $p \in \mathbb{Z}[t, t^{-1}]$ can be expressed uniquely in the form $\sum_{i=m}^M p_i t^i$ where m, M and all the $p_i \in \mathbb{Z}$ and neither p_m nor p_M is 0. Denote by $m(p)$ and $M(p)$ the values of m and M in this expression and define the *length* of p , $\|p\|$, to be $\sum_{i=m}^M |p_i|$. For $\lambda \in \Lambda_R$, define $\bar{\lambda}$ to be the image of λ under the unique nontrivial (ring) involution of Λ_R . Correspondingly, for $p \in \mathbb{Z}[t, t^{-1}]$, define $\bar{p}$ to be the image of p under the corresponding involution of $\mathbb{Z}[t, t^{-1}]$, which takes t to $\pm 1/t$, the sign depending on the determinant of R . This choice of sign serves to guarantee that $\alpha_R(\bar{p}) = \overline{\alpha_R(p)}$.

The action by R gives rise to the structure of a $\mathbb{Z}[t, t^{-1}]$ -module on $\mathbb{Z}^2 = \langle x, y \mid [x, y] \rangle$, where t acts by R . (We will continue to denote these generators of $\mathbb{Z}^2$ by x and y .) This induces a Λ_R -action. Let M_R refer to $\mathbb{Z}^2$ with this Λ_R -action.

The map R , being hyperbolic, has two real eigenvalues, τ and $\pm 1/\tau$, corresponding to two eigenvectors $\mathbf{v}_e(R)$ and $\mathbf{v}_c(R)$, taken to be expanding and contracting respectively. (Clearly, each of these eigenvectors is defined only up to a multiplicative constant.) The line spanned by $\mathbf{v}_c(R)$ will be called the *contracting line* and the line spanned by $\mathbf{v}_e(R)$ the *expanding line*. For any $z \in M_R$, let $d_c(z)$ be the (Euclidean) distance of z from the contracting line. Then the following diagram commutes:

$$\begin{array}{ccc}
 M_R & \xrightarrow{R} & M_R \\
 d_c \downarrow & & d_c \downarrow \\
 \mathbb{R}_{\geq 0} & \xrightarrow{\cdot |\tau|} & \mathbb{R}_{\geq 0}.
 \end{array}$$

Similarly, let $d_e(z)$ be the distance of z from the expanding line, so that the following diagram commutes:

$$\begin{array}{ccc} M_R & \xrightarrow{R \cdot} & M_R \\ d_e \downarrow & & d_e \downarrow \\ \mathbb{R}_{\geq 0} & \xrightarrow{\div |\tau|} & \mathbb{R}_{\geq 0}. \end{array}$$

Lemma 23. *Let $\Lambda_R = \mathbb{Z}[t]/p_R(t)$, where R is a hyperbolic automorphism of $\mathbb{Z}^2$. Then there exist $C_1, C_2 > 0$ with the following property. Let $\lambda \in \Lambda_R$ and let α_R represent as usual the natural projection of $\mathbb{Z}[t, t^{-1}]$ to Λ_R . Let $p, q \in \alpha_R^{-1}(\lambda)$, with p a minimal-length element of this set. Then, for any $z \in M_R$,*

$$d_c(\lambda z) < |\tau|^{M(q)} (C_1(\|q\| - \|p\|) + C_2)d_c(z)$$

and

$$d_c(\bar{\lambda}z) < |\tau|^{-m(q)} (C_1(\|q\| - \|p\|) + C_2)d_c(z).$$

Furthermore,

$$d_e(\lambda z) < |\tau|^{-m(q)} (C_1(\|q\| - \|p\|) + C_2)d_e(z)$$

and

$$d_e(\bar{\lambda}z) < |\tau|^{M(q)} (C_1(\|q\| - \|p\|) + C_2)d_e(z)$$

Proof. We prove only the first statement; the second follows by applying the involution $\bar{}$ throughout the proof and the third and fourth then follow by replacing R with R^{-1} (note that $|\text{tr } R^{-1}| = |\text{tr } R|$ since $|\det R| = 1$).

Letting brackets represent the greatest integer function, we have D depending only on R such that

$$\sum_{i=-\infty}^{\infty} [Dq_i] \leq \|q\| - \|p\|,$$

for otherwise we could apply some multiple of the equation $t^2 \pm 1 = (\text{tr } R)t$ to q more than $\|q\| - \|p\|$ times, shortening q each time without changing $\alpha_R(q)$. This would be a contradiction since p is of minimal length in $\alpha_R^{-1}(\lambda)$. If $|\text{tr } R| = 1$ or 2 (in which case the constant term has negative sign, since R is hyperbolic), we must instead use the equations $3 = t^2 + t^{-2}$ and $6 = t^2 + t^{-2}$ respectively. Thus, for any hyperbolic R ,

$$\begin{aligned} \frac{d_c(\lambda z)}{d_c(z)} &\leq \frac{\|q\| - \|p\| + 1}{D} \tau^{M(q)} + \sum_{i=0}^{\infty} \frac{1}{D} |\tau|^{M(q)-i-1} \\ &= \frac{\|q\| - \|p\|}{D} |\tau|^{M(q)} + \frac{1}{D} \sum_{i=0}^{\infty} |\tau|^{M(q)-i} \\ &= |\tau|^{M(q)} \left(\frac{\|q\| - \|p\|}{D} + \frac{1}{D} \frac{1}{1 - |\tau|^{-1}} \right), \end{aligned}$$

as claimed. $\square$

If $p_1, p_2 \in \mathbb{Z}[t, t^{-1}]$, define the *length* of (p_1, p_2) to be $\|p_1\| + \|p_2\|$. We mean by $M(p_1, p_2)$ the greater of $M(p_1)$ and $M(p_2)$ and by $m(p_1, p_2)$ the lesser of $m(p_1)$ and $m(p_2)$.

Observe that conjugation by c , denoted by ρ , sends $a^i b^j$ to $a^{i'} b^{j'}$, where $(i', j') = R(i, j)$. Let $\beta: G'_R \leftrightarrow M_R$ send a to x and b to y . Then $\beta(a^c) = R(\beta(a))$ and $\beta(b^c) = R(\beta(b))$, so this diagram commutes:

$$\begin{array}{ccc} G'_R & \xrightarrow{\cdot c} & G'_R \\ \beta \downarrow & & \beta \downarrow \\ M_R & \xrightarrow{R \cdot} & M_R. \end{array}$$

Furthermore, β induces an obvious one-to-one length-preserving correspondence θ between N_{ab} and $\mathbb{Z}[t, t^{-1}]^2$ such that this diagram commutes:

$$\begin{array}{ccc} N_{ab} & \xrightarrow{\theta} & \mathbb{Z}[t, t^{-1}]^2 \\ \pi \downarrow & & \downarrow (p_1, p_2) \mapsto p_1 x + p_2 y \\ G'_R & \xrightarrow{\beta} & M_R. \end{array}$$

Definition 4. If $v = \sum_{j=1}^n u_{k_j}^{i_j}$ is a word in $\mathbb{Z}^T$, let $M(v)$ and $m(v)$ be the maximum and minimum of the i_j .

Proposition 20 will follow from Proposition 21 and from the following

Proposition 24. Let $G_R = \mathbb{Z}^2 \rtimes_R \langle c \rangle$, where R is a hyperbolic automorphism of $\mathbb{Z}^2$. Then there are $D_2 > 1$ and $D_1 > D_2 \ln |\tau| / 4$ with the following property. Let $g = uc^z \in G_R$, where $u \in G'_R$. Let $v, v' \in N_{ab}$ be elements of $\pi^{-1}(u)$, with v of minimal length. (We remind the reader that π is the projection from N_{ab} to G'_R .) Then $|\tau|^{2M(v)-2M(v')} < D_1(l(v') - l(v)) + D_2$ and $|\tau|^{2m(v')-2m(v)} < D_1(l(v') - l(v)) + D_2$.

The proof follows easily from the following proposition about M_R (whose proof we postpone) and from Lemma 23.

Proposition 25. Let M_R denote $\mathbb{Z}^2$ equipped with an action by $\mathbb{Z}[t, t^{-1}]$, where t acts by some hyperbolic automorphism R of $\mathbb{Z}^2$. Let x and y denote the standard generators of M_R as an abelian group. Then there exists $e > 0$ with the following property. Let $z \in M_R$ and $p_1, p_2 \in \mathbb{Z}[t, t^{-1}]$ such that $p_1 x + p_2 y = z$ and $\|p_1\| + \|p_2\|$ is minimal subject to this condition. Then $d_c(z) = d_c(p_1 x + p_2 y) > e |\tau|^{M(p_1, p_2)}$ and $d_c(\overline{p_1} x + \overline{p_2} y) > e |\tau|^{-m(p_1, p_2)}$.

Proof of Proposition 24. Let $(p_1, p_2) = \theta(v)$ and let $z = \beta(u) = p_1 x + p_2 y$. Then (p_1, p_2) is in fact of minimal length such that $p_1 x + p_2 y = z$ (since θ is length-preserving and v is a minimal-length element of $\pi^{-1}(u)$), so by Proposition 25 $e |\tau|^{M(p_1, p_2)} < d_c(z)$ and $e |\tau|^{-m(p_1, p_2)} < d_c(\overline{p_1} x + \overline{p_2} y)$.

If similarly $(p'_1, p'_2) = \theta(v')$ then $p'_1x + p'_2y = \beta(\pi(v')) = \beta(u) = z$. Then by Lemma 23 we have

$$\begin{aligned} e|\tau|^{M(p_1, p_2)} &< d_c(z) \leq d_c(p_1x) + d_c(p_2y) \\ &< |\tau|^{M(p'_1, p'_2)} (C_1(\|p'_1\| + \|p'_2\| - \|p_1\| - \|p_2\|) + 2C_2) \\ &\quad \cdot \max(d_c(x), d_c(y)) \end{aligned}$$

and

$$\begin{aligned} e|\tau|^{-m(p_1, p_2)} &< d_c(\overline{p_1}x + \overline{p_2}y) \leq d_c(\overline{p_1}x) + d_c(\overline{p_2}y) \\ &< |\tau|^{-m(p'_1, p'_2)} (C_1(\|p'_1\| + \|p'_2\| - \|p_1\| - \|p_2\|) + 2C_2) \\ &\quad \cdot \max(d_c(x), d_c(y)), \end{aligned}$$

where C_1 and C_2 depend only on R as in Lemma 23. Thus

$$\begin{aligned} |\tau|^{M(p_1, p_2) - M(p'_1, p'_2)} &< (C_1(\|p'_1\| + \|p'_2\| - \|p_1\| - \|p_2\|) + 2C_2) \max(d_c(x), d_c(y)) \end{aligned}$$

and

$$\begin{aligned} |\tau|^{m(p'_1, p'_2) - m(p_1, p_2)} &< (C_1(\|p'_1\| + \|p'_2\| - \|p_1\| - \|p_2\|) + 2C_2) \max(d_c(x), d_c(y)). \end{aligned}$$

Clearly, $M(v) = M(p_1, p_2)$, $m(v) = m(p_1, p_2)$ and $l(v) = \|p_1\| + \|p_2\|$, and similarly for v' and (p'_1, p'_2) . This gives us that

$$\begin{aligned} |\tau|^{M(v) - M(v')} &= |\tau|^{M(p_1, p_2) - M(p'_1, p'_2)} \\ &< (C_1(\|p'_1\| + \|p'_2\| - \|p_1\| - \|p_2\|) + 2C_2) \max(d_c(x), d_c(y)) \\ &= (C_1(l(v') - l(v)) + 2C_2) \max(d_c(x), d_c(y)) \end{aligned}$$

and similarly

$$|\tau|^{m(v') - m(v)} < (C_1(l(v') - l(v)) + 2C_2) \max(d_c(x), d_c(y)).$$

(If the D_2 resulting from the above expressions is not greater than 1, we may clearly increase it without destroying the result. We may similarly increase D_1 to be greater than $D_2 \ln |\tau| / 4$.) $\square$

Proof of Proposition 20. Let $w \in F_S$ be a minimal-length element of $\phi^{-1}(v) \cap \sigma_R^{-1}(g)$ and let $w' \in F_S$ be a minimal-length element of $\sigma_R^{-1}(g)$. By definition, $|g| = l(w')$.

The bound $l(w') \leq l(w)$ is obvious, so we need only prove $l(w) - D \leq l(w')$. Let $v' = \phi(w')$. We note that $\pi(v') = u = \pi(v)$. It is clear from the definition of M and m that $M(v) \geq m(v)$, so we cannot have both $M(v)$ negative and $m(v)$ positive, and similarly for $M(v')$ and $m(v')$. We may thus assume without loss of generality that $m(v)$ and $m(v')$ are both nonpositive. we will further assume that $M(v)$ and

$M(v')$ are both nonnegative; the other cases are easier and are left to the reader. In this case, we have by Lemma 21

$$\begin{aligned} [l(w) - l(w')] - [l(v) - l(v')] &\leq 2[M(v) - M(v')] + 2[m(v') - m(v)] \\ &\quad + \min(|z - M(v)| - M(v), |z - m(v)| + m(v)) \\ &\quad - \min(|z - M(v')| - M(v'), |z - m(v')| + m(v')). \end{aligned}$$

Noting that $|z - M(v)| - M(v) \geq |z - m(v)| + m(v)$ if and only if $|z - M(v')| - M(v') \geq |z - m(v')| + m(v')$ if and only if $z \geq 0$, we find using the triangle inequality that

$$\begin{aligned} [l(w) - l(w')] - [l(v) - l(v')] - 2[M(v) - M(v')] - 2[m(v') - m(v)] &\leq \max(|z - M(v)| - |z - M(v')| + M(v') - M(v), \\ &\quad |z - m(v)| - |z - m(v')| + m(v) - m(v')) \\ &\leq \max(2|M(v') - M(v)|, 2|m(v) - m(v')|) \\ &\leq 2\max(M(v) - M(v'), 0) + 2\max(m(v') - m(v), 0). \end{aligned}$$

By Proposition 24 and using that $l(v') \geq l(v)$ since v is given to be of minimal length in $\pi^{-1}(u)$, we get

$$\begin{aligned} [l(w) - l(w')] - [l(v) - l(v')] &\leq 4\max(M(v) - M(v'), 0) + 4\max(m(v') - m(v), 0) \\ &< 2\max(\log_{|\tau|}[D_1(l(v') - l(v)) + D_2], 0) + 2\max(\log_{|\tau|}[D_1(l(v') - l(v)) + D_2], 0) \\ &= 4\log_{|\tau|}[D_1(l(v') - l(v)) + D_2]. \end{aligned}$$

Rearranging yields that $l(w) - l(w') \leq 4\log_{|\tau|}[D_1(l(v') - l(v)) + D_2] - (l(v') - l(v))$.

It now remains to show that $f(x) = 4\log_{|\tau|}(D_1x + D_2) - x$ attains an absolute maximum value on the domain $[0, \infty)$, which can be done by differential calculus. (This absolute maximum value will depend only on R since τ , D_1 and D_2 depend only on R .) We compute $f'(x) = 4D_1/[(D_1x + D_2)\ln|\tau|] - 1$ and observe that it is a strictly decreasing function on the domain of interest, so (by the first derivative test for absolute maxima), if it has a zero, the absolute maximum of f is attained there. We find that $\lim_{x \rightarrow \infty} f'(x) = -1$ while $f'(0) = 4D_1/(D_2 \ln|\tau|) - 1 > 0$ since we took $D_1 > D_2 \ln|\tau|/4$. Since f' is continuous on $[0, \infty)$, we are done. $\square$

The proof of Proposition 20 thus reduces to that of Proposition 25. This in turn follows from the following two results.

Lemma 26. *Let M_R denote $\mathbb{Z}^2$ equipped with an action by $\mathbb{Z}[t, t^{-1}]$, where t acts by some hyperbolic automorphism R of $\mathbb{Z}^2$. Let x and y denote the standard generators of M_R as an abelian group. Let Λ_R denote the quotient of $\mathbb{Z}[t, t^{-1}]$ by the characteristic polynomial of R . Let $\lambda_1, \lambda_2 \in \Lambda_R$ and $p_1, p_2 \in \mathbb{Z}[t, t^{-1}]$ be minimal-length elements of $\alpha_R^{-1}(\lambda_1)$ and $\alpha_R^{-1}(\lambda_2)$ respectively. Then*

$$6(d_c(x) + d_c(y)) |\tau|^{M(p_1, p_2)+1} > d_c(p_1x + p_2y)$$

and

$$6(d_c(x) + d_c(y)) |\tau|^{-m(p_1, p_2)+1} > d_c(\overline{p_1}x + \overline{p_2}y).$$

Proof. We prove the first statement of the lemma; the second will follow by applying the involution $\overline{}$ throughout the proof.

We know, assuming $|\operatorname{tr} R| > 2$, that no coefficient in p_1 or p_2 can have absolute value greater than $|\operatorname{tr} R| + 2$, since otherwise we could apply the equation $\pm t^2 - (\operatorname{tr} R)t \pm 1 = 0$ to produce a shorter element of $\alpha_R^{-1}(\lambda_1)$ or $\alpha_R^{-1}(\lambda_2)$. If $|\operatorname{tr} R| = 1$ or 2 the same result holds, since instead we use the equation $t^2 + t^{-2} = 3$ or 6 , respectively. Thus, in either case, given $z \in M_R$, both $d_c(p_1 z)/d_c(z)$ and $d_c(p_2 z)/d_c(z)$ are

$$\leq \sum_{i=-\infty}^{M(p_1, p_2)} (|\operatorname{tr} R| + 2) |\tau|^i = (|\operatorname{tr} R| + 2) \frac{|\tau|^{M(p_1, p_2)+1}}{|\tau| - 1}.$$

If we denote $|\operatorname{tr} R|$ by r then this equals

$$\frac{2(r+2) |\tau|^{M(p_1, p_2)+1}}{r + \sqrt{r^2 \mp 4} - 2} < 6 |\tau|^{M(p_1, p_2)+1}$$

since either $r > 2$ or the sign inside the square root is positive. It follows that

$$d_c(p_1 x + p_2 y) \leq d_c(p_1 x) + d_c(p_2 y) < 6(d_c(x) + d_c(y)) |\tau|^{M(p_1, p_2)+1},$$

as claimed. $\square$

Proposition 27. *Let M_R denote $\mathbb{Z}^2$ equipped with an action by $\mathbb{Z}[t, t^{-1}]$, where t acts by some hyperbolic automorphism R of $\mathbb{Z}^2$. Let x and y denote the standard generators of M_R as an abelian group and let P and Q be positive numbers. Then there exists a natural number n with the following property. Let $z \in M_R$ and (p_1, p_2) be a minimal-length element of $\mathbb{Z}[t, t^{-1}]^2$ such that $p_1 x + p_2 y = z$. Then $(Pd_c(x) + Qd_c(y)) |\tau|^{M(p_1, p_2)-n} < d_c(z)$.*

We postpone the proof and instead first give the

Proof of Proposition 25. As in the proof of Lemma 26, we prove the first statement of the lemma; the second will follow by applying the involution $\overline{}$ throughout the proof.

If $p \in \mathbb{Z}[t, t^{-1}]$, we let $\operatorname{head}_m(p)$ denote the sum of those terms of p of degree at least m , and $\operatorname{tail}_m(p)$ the sum of all remaining terms. (Thus $p = \operatorname{head}_m(p) + \operatorname{tail}_m(p)$.) If m is taken to be $M(p_1, p_2) - n$, with n given by Proposition 27 with $P = 7$ and $Q = 6$, we have $d_c(\operatorname{head}_m(p_1)x + \operatorname{head}_m(p_2)y) > (7d_c(x) + 6d_c(y)) |\tau|^{M(p_1, p_2)-n}$ by that proposition. However, $M(\operatorname{tail}_m(p_1), \operatorname{tail}_m(p_2)) = m - 1$. Thus, by Lemma 26, we have

$$\begin{aligned} d_c(\operatorname{tail}_m(p_1)x + \operatorname{tail}_m(p_2)y) &< 6(d_c(x) + d_c(y)) |\tau|^m \\ &= 6(d_c(x) + d_c(y)) |\tau|^{M(p_1, p_2)-n}, \end{aligned}$$

since certainly if (p_1, p_2) is of minimal length such that $p_1 x + p_2 y = z$ then each of the p_i is of minimal length in $\alpha_R^{-1}(\alpha_R(p_i))$. (We again remind the reader that α_R is the projection from $\mathbb{Z}[t, t^{-1}]$ to M_R .) It follows that

$$\begin{aligned} d_c(z) &\geq d_c(\operatorname{head}_m(p_1)x + \operatorname{head}_m(p_2)y) - d_c(\operatorname{tail}_m(p_1)x + \operatorname{tail}_m(p_2)y) \\ &> d_c(x) |\tau|^{M(p_1, p_2)-n}. \end{aligned}$$

We may thus take e to be $d_c(x) |\tau|^{-n}$. $\square$

It remains now (for the proof of Proposition 20) to prove Proposition 27. It will follow from the following three results.

Lemma 28. *Let M_R denote $\mathbb{Z}^2$ equipped with an action by $\mathbb{Z}[t, t^{-1}]$, where t acts by some hyperbolic automorphism R of $\mathbb{Z}^2$. Let x and y denote the standard generators of M_R as an abelian group and let $d \in \mathbb{R}$. Then there exist $C_1 > 0$ and $C_2 \in \mathbb{R}$ with the following property. Let $p_1, p_2 \in \mathbb{Z}[t, t^{-1}]$ be such that $d_c(p_1x + p_2y) \leq d$ and the first coordinate of $p_1x + p_2y \in M_R$ is w . Assume that $m(p_1, p_2) \geq 0$. Then $\|p_1\| + \|p_2\| \geq C_1|w| + C_2$.*

Proof. Since $m(p_1, p_2) \geq 0$, $d_e(p_1x + p_2y) \leq \|p_1\| + \|p_2\|$, since $p_1x + p_2y$ is the sum of $\|p_1\| + \|p_2\|$ vectors v_i ($= t^j x$ or $t^j y$, $j \in \mathbb{Z}$) such that $d_e(v_i) \leq 1$. However, the nearest point to $p_1x + p_2y$ on the contracting line has first coordinate of absolute value at least $|w| - d$. This point is thus at distance at least $C_1(|w| - d)$ from the expanding line, where C_1 is a positive constant depending only on R . Since $p_1x + p_2y$ is within distance d of this point, $d_e(p_1x + p_2y) \geq C_1(|w| - d) - d$, as claimed. The result follows by transitivity if we let $C_2 = -d(C_1 + 1)$. $\square$

We postpone the proof of the next two results.

Proposition 29. *Let M_R denote $\mathbb{Z}^2$ equipped with an action by $\mathbb{Z}[t, t^{-1}]$, where t acts by some hyperbolic automorphism R of $\mathbb{Z}^2$. Let x and y denote the standard generators of M_R as an abelian group and let $d \geq 0$. Then there exist D_1, D_2 and D_3 such that the following holds. Let $z \in M_R$ be such that $d_c(z) \leq d$ and the first coordinate of z is w . Then there exist $p_1, p_2 \in \mathbb{Z}[t, t^{-1}]$ such that $p_1x + p_2y = z$ and $\|p_1\| + \|p_2\| \leq D_1 + \max(0, D_2 \ln(D_3 |w|))$.*

Proposition 30. *Let M_R denote $\mathbb{Z}^2$ equipped with an action by $\mathbb{Z}[t, t^{-1}]$, where t acts by some hyperbolic automorphism R of $\mathbb{Z}^2$. Let x and y denote the standard generators of M_R as an abelian group and let $z \in M_R$. Let P be the set of all $(p_1, p_2) \in \mathbb{Z}[t, t^{-1}]^2$ such that $p_1x + p_2y = z$. Then P contains only finitely many elements of minimal length.*

Proof of Proposition 27. It will suffice to show for some n that, if $M(p_1, p_2) \geq n$, then $Pd_c(x) + Qd_c(y) < d_c(z)$.

Suppose the conclusion does not hold for z, p_1 and p_2 ; that is, suppose that $Pd_c(x) + Qd_c(y) \geq d_c(z)$ where $z = p_1x + p_2y$. Then applying Lemma 28 with $d = Pd_c(x) + Qd_c(y)$ (which, it should be noted, depends only on R, P and Q) yields $C_1 > 0$ and C_2 such that

$$\|p_1\| + \|p_2\| \geq C_1|w| + C_2,$$

where w is the first coordinate of z . However, since p_1 and p_2 were taken so that $\|p_1\| + \|p_2\|$ would be minimal subject to the condition $p_1x + p_2y = z$, applying Proposition 29 with the same value of d gives D_1, D_2 and D_3 such that $\|p_1\| + \|p_2\| \leq D_1 + \max(0, D_2 \ln(D_3 |w|))$. Thus

$$D_1 + \max(0, D_2 \ln(D_3 |w|)) \geq C_1|w| + C_2.$$

Since the left-hand side depends logarithmically on $|w|$ while the right-hand side is increasing and depends linearly on it, there exists w' dependent only on R, P and Q (via D_1, D_2, D_3, C_1 and C_2) such that $w < w'$. Thus z is a lattice point constrained to lie within a compact set, since it is confined to the intersection of finite-width bands about the y -axis and the contracting line. There are only finitely

many such points. Thus there are also only finitely many possibilities for p_1 and p_2 , by Proposition 30. The result follows by taking n to be 1 more than the maximal $M(p_1, p_2)$ for any of these finitely many possibilities. $\square$

Proposition 29 will follow from the following

Lemma 31. *Let R be a hyperbolic automorphism of $\mathbb{Z}^2$. Let S be the set of all integers appearing in the first row of the matrix for some nonnegative power of R . Then there exist C_1 , C_2 and C_3 such that any $n \in \mathbb{Z}$ can be expressed as a word in the elements of S of length at most $C_1 + \max(0, C_2 \ln(C_3 |n|))$.*

Proof. The matrix for R consists of two column vectors, which we denote (p, q) and (r, s) . Each can be decomposed as the sum of a (real) vector in the expanding line and one in the contracting line; thus $(p, q) = (p_e, q_e) + (p_c, q_c)$ and $(r, s) = (r_e, s_e) + (r_c, s_c)$ where (p_e, q_e) and (r_e, s_e) lie in the expanding line and (p_c, q_c) and (r_c, s_c) in the contracting line. If we denote the expanding eigenvalue of R by τ then we find that

$$S = \left\{ p_m = p_e \tau^m \pm \frac{p_z}{\tau^m}, r_m = r_e \tau^m \pm \frac{r_z}{\tau^m} \mid m \geq 0 \right\},$$

where the $\pm$ signs depend on the determinant of R and the parity of m .

Let C_1 be the maximal length of any minimal-length word in the elements of S representing a number of absolute value $\leq |p_e|$. Let C_2 be $|\tau| (1 + |p_z|) / \ln |\tau|$ and C_3 be $|\tau/p_e|$.

Consider $n, m \in \mathbb{Z}$ with $|n| \leq |p_e \tau^m|, m > 0$. Then there must exist $k \in \mathbb{Z}$ with $|k| < |\tau|$ such that $|n + k p_e \tau^{m-1}| \leq |p_e \tau^{m-1}|$. It follows that

$$|n + k p_{m-1}| \leq |n + k p_e \tau^{m-1}| + \left| \frac{k p_c}{\tau^{m-1}} \right| \leq |p_e \tau^{m-1}| + |k p_c| \leq |p_e \tau^{m-1}| + |k p_c|,$$

so there exists $l \in \mathbb{Z}$ with $|l| \leq |k p_c| < |\tau p_c|$ such that $|n + k p_{m-1} + l p_0| \leq |p_e \tau^{m-1}|$. (Note that $p_0 = 1$.) The result follows by induction on m . $\square$

Proof of Proposition 29. By Lemma 31 applied to the hyperbolic automorphism R^{-1} , there exist C_1 , C_2 and C_3 depending only on R and $p_1, p'_2 \in \mathbb{Z}[t, t^{-1}]$ such that $p_1 x + p'_2 y$ has first coordinate w and

$$\|p_1\| + \|p'_2\| \leq C_1 + \max(0, C_2 \ln(C_3 |w|)).$$

Furthermore, $M(p_1, p'_2) \leq 0$. Then

$$d_c(p_1 x + p'_2 y) \leq C_1 + \max(0, C_2 \ln(C_3 |w|))$$

since $p_1 x + p'_2 y$ is the sum of $C_1 + \max(0, C_2 \ln(C_3 |w|))$ vectors $v_i (= t^j x$ or $t^j y$, $j \in \mathbb{Z})$ such that $d_c(v_i) \leq 1$. Thus $p_1 x + p'_2 y$ is within $\sqrt{2}[C_1 + \max(0, C_2 \ln(C_3 |w|))]$ vertically of the point on the contracting line with first coordinate w , so within $\sqrt{2}[C_1 + \max(0, C_2 \ln(C_3 |w|)) + d]$ of z . (It is here that we use the condition that $|zeta| \leq 1$.) Defining p_2 to be

$$p'_2 + (\text{some appropriate constant term}),$$

we see that $D_1 = (1 + \sqrt{2})C_1 + \sqrt{2}d$, $D_2 = (1 + \sqrt{2})C_2$ and $D_3 = C_3$ give what we seek. $\square$

The proof of Proposition 20 has now been reduced to that of Proposition 30, which follows from the following

Lemma 32. *Let M_R denote $\mathbb{Z}^2$ equipped with an action by $\mathbb{Z}[t, t^{-1}]$, where t acts by some hyperbolic automorphism R of $\mathbb{Z}^2$. Let x and y denote the standard generators of M_R as an abelian group and let $z \neq \mathbf{0} \in M_R$ and $l \in \mathbb{N}$. Then there exists an integer $N = N(z, l, R)$ with the following property. Let $p_1, p_2 \in \mathbb{Z}[t, t^{-1}]$ such that*

$$p_1x + p_2y = z$$

and $\|p_1\| + \|p_2\| = l$. Then at least one of p_1 and p_2 must have a nonzero term whose degree has absolute value less than N .

Proof. Express z (viewed as an element of $\mathbb{Z}^2 \subset \mathbb{R}^2$) as $z = z_e \mathbf{v}_e(R) + z_c \mathbf{v}_c(R)$, $z_e, z_c \in \mathbb{R}$. (We remind the reader that $\mathbf{c}_e(R)$ and $\mathbf{v}_c(R)$ generate the expanding and contracting eigenspaces respectively.) Since both eigenvalues of R are irrational, no nonzero real multiple of $\mathbf{v}_c(R)$ lies in the $\mathbb{Z}^2$ lattice. In particular, since $z \neq \mathbf{0}$, $z_c \mathbf{v}_c(R) \notin \mathbb{Z}^2$. Let $D > 0$ be the (Euclidean Hausdorff) distance between $z_c \mathbf{v}_c(R)$ and $\mathbb{Z}^2$, and let $N > \log_{|\tau|}(2l/D)$.

If $p \in \mathbb{Z}[t, t^{-1}]$, let $p_{\geq m} \in \mathbb{Z}[t, t^{-1}]$ denote the sum of all terms of p of degree $\geq m$, and analogously for $p_{\leq m}$. Suppose (for a contradiction) $p_1 = p_{1 \geq N} + p_{1 \leq -N}$ and similarly for p_2 . Then

$$p_1x + p_2y = p_{1 \geq N}x + p_{2 \geq N}y + p_{1 \leq -N}x + p_{2 \leq -N}y.$$

We define $z_{e \geq N}$ and $z_{c \geq N} \in \mathbb{R}$ such that

$$p_{1 \geq N}x + p_{2 \geq N}y = z_{e \geq N} \mathbf{v}_e(R) + z_{c \geq N} \mathbf{c}(R),$$

and similarly for $z_{e \leq -N}$ and $z_{c \leq -N}$. Then each term $a_i t^i x$ or $a_i t^i y$ of $p_{1 \geq N}x + p_{2 \geq N}y$ contributes at most $|a_i|/\tau^N$ of Euclidean length to $z_{c \geq N} \mathbf{v}_c(R)$. Since the sum of all the a_i is at most l , $z_{c \geq N} \mathbf{v}_c(R)$ has Euclidean length at most l/τ^N . Similarly, $z_{e \leq -N} \mathbf{v}_e(R)$ has Euclidean length at most l/τ^N . Then

$$p_{1 \leq -N}x + p_{2 \leq -N}y = z_c \mathbf{v}_c(R) - z_{c \geq N} \mathbf{v}_c(R) + z_{e \leq -N} \mathbf{v}_e(R) \in \mathbb{Z}^2$$

lies within Euclidean distance $2l/\tau^N < D$ of $z_c \mathbf{v}_c(R)$. But this is impossible by the definition of D . $\square$

Proof of Proposition 30. The result is trivial for $z = \mathbf{0}$, since then $P = \{(0, 0)\}$, so we assume $z \neq \mathbf{0}$.

Suppose P contains infinitely many minimal-length elements. Let l be the length of these minimal-length elements. Then they each contain some term in which t is raised to an exponent of absolute value $< N = N(z, l, R)$, by Lemma 32. Thus each minimal-length element of P comes from a length- $(l-1)$ minimal-length expression for one of the finitely many possibilities $z \pm t^{-N+1}x$, $z \pm t^{-N+1}y$, $z \pm t^{-N+2}x$, $\dots$, $z \pm t^{N-1}y$. The result follows by induction on l . $\square$

This concludes the proof of Proposition 20. We now need only to prove Proposition 22.

Lemma 33. *Let $G_R = \mathbb{Z}^2 \rtimes_R \mathbb{Z}$, where R is a hyperbolic automorphism of $\mathbb{Z}^2$, and let $z = (z_1, z_2) \in G'_R$ be such that $|z_1|, |z_2| < |\text{tr } R|^m$. Then there exists a word $w \in \sigma_R^{-1}(\beta^{-1}(z))$ such that $l(w) \leq 2^{m+1} + 4m - 1$. (Recall that β is the correspondence between G'_R and $\mathbb{Z}^2$ and σ_R is the surjection from F_S to G_R .) If $|\text{tr } R| = 1$ then if $|z_1|, |z_2| < 3^m$ then there exists $w \in \sigma_R^{-1}(\beta^{-1}(z))$ such that $l(w) \leq 2^{m+1} + 8m - 1$. If $|\text{tr } R| = 2$ then if $|z_1|, |z_2| < 6^m$ then there exists $w \in \sigma_R^{-1}(\beta^{-1}(z))$ such that $l(w) \leq 2^{m+1} + 8m - 1$.*

Proof. We begin with the proof of the first claim.

Since $|z_1|, |z_2| < |\operatorname{tr} R|^m$, we can express them in the form

$$z_1 = \sum_{i=0}^m z_{1i} |\operatorname{tr} R|^i,$$

$$z_2 = \sum_{i=0}^m z_{2i} |\operatorname{tr} R|^i,$$

where all the $|z_{1i}|$ and $|z_{2i}| \leq |\operatorname{tr} R|/2$. But since, in Λ_R , $|\operatorname{tr} R| = \pm\tau \pm \tau^{-1}$, $|\operatorname{tr} R|^i = (\pm\tau \pm \tau^{-1})^i = \sum_{j=0}^i \pm \binom{i}{j} \tau^{2j-i}$. Thus we can reexpress z_1 as $\sum_{k=-m}^m z'_{1k} \tau^k$, where

$$\sum_{k=-m}^m |z'_{1k}| \leq \sum_{i=0}^m \sum_{j=0}^i \binom{i}{j} = \sum_{i=0}^m 2^i = 2^{m+1} - 1,$$

and similarly for z_2 . We thus have $z = (\sum_{k=-m}^m z'_{1k} t^k)x + (\sum_{k=-m}^m z'_{2k} t^k)y$. Thus there exists $v = \sum_{j=1}^n u_{k_j}^{c_{i_j}} \in \mathbb{Z}^T$ with $\pi(v) = \beta^{-1}(z)$ and all $k_j = 1$ or 2 such that all i_j satisfy $-m \leq i_j \leq m$ and $n \leq 2^{m+1} - 1$. By Lemma 21, this gives a $w \in \sigma_R^{-1}(\beta^{-1}(z))$ such that $l(w) \leq 2^{m+1} + 4m - 1$, as claimed.

The proofs of the second and third claims are analogous; just replace $|\operatorname{tr} R|$ by 3 and 6 respectively and the equation $|\operatorname{tr} R| = \pm\tau \pm \tau^{-1}$ by the equations $3 = \tau^2 + \tau^{-2}$ and $6 = \tau^2 + \tau^{-2}$ respectively. $\square$

Proof of Proposition 22. We first construct $B_{m,n}$. Recall that β is the correspondence between G'_R and M_R and θ is that between N_{ab} and $\mathbb{Z}[t, t^{-1}]^2$. Let $u \in G'_R$ and set $v \in N_{ab}$ to be a minimal-length element of $\pi^{-1}(u)$. Then by Lemma 23 (setting $(q_1, q_2) = (p_1, p_2) = \theta(v)$)

$$\begin{aligned} d_c(\beta(u)) &\leq d_c(p_1x) + d_c(p_2y) \\ &< |\tau|^{M(\theta(v))} (C_1(\|q_1\| + \|q_2\| - \|p_1\| - \|p_2\|) + 2C_2) \max(d_c(x), d_c(y)) \\ &= 2C_2 |\tau|^{M(\theta(v))} \max(d_c(x), d_c(y)) \end{aligned}$$

and

$$\begin{aligned} d_e(\beta(u)) &< |\tau|^{-m(\theta(v))} (C_1(\|q_1\| + \|q_2\| - \|p_1\| - \|p_2\|) + 2C_2) \\ &\quad \cdot \max(d_e(x), d_e(y)) \\ &= 2C_2 |\tau|^{-m(\theta(v))} \max(d_e(x), d_e(y)) \end{aligned}$$

where C_2 depends only on R . Then if we assume

$$d_c(\beta(u)) > 2C_2 |\tau|^n \max(d_c(x), d_c(y)),$$

and

$$d_e(\beta(u)) > 2C_2 |\tau|^n \max(d_e(x), d_e(y)),$$

this implies that $M(\theta(v)) > n$, $m(\theta(v)) < -n$, so $M(v) > n$, $m(v) < -n$. We let $B_{m,n}$ be the set of $u = a^i b^j \in [G_R, G_R]$ such that $|i|, |j| < |\operatorname{tr} R|^m$,

$$d_c(\beta(a^i b^j)) > 2C_2 |\tau|^n \max(d_c(x), d_c(y))$$

and

$$d_e(\beta(a^i b^j)) > 2C_2 |\tau|^n \max(d_e(x), d_e(y)).$$

If $|\operatorname{tr} R| = 1$ then replace $|\operatorname{tr} R|$ in the preceding sentence by 3.

We proceed with the proof of the rest of the proposition, first assuming $|\operatorname{tr} R| > 2$.

Let K and L be positive integers such that

$$(L + 2C_2 |\tau|^n) \frac{\max(d_c(x), d_c(y), d_e(x), d_e(y))}{\min(d_c(x), d_e(x))} < K < |\operatorname{tr} R|^m - L;$$

such K and L clearly exist for sufficiently large m and we can arrange for L to grow proportionally to $|\operatorname{tr} R|^m$ for R and n fixed. Let $z = Kx$. Addition or subtraction of x or y does not change $d_c(z)$ by more than $\max(d_c(x), d_c(y))$ nor $d_e(z)$ by more than $\max(d_e(x), d_e(y))$. Thus any element $z' = (z'_1, z'_2) \in \mathbb{Z}^2$ at $\{x, y\}$ -distance less than L from z satisfies

$$d_c(z') > K d_c(x) - L \max(d_c(x), d_c(y)) > 2C_2 |\tau|^n \max(d_c(x), d_c(y)),$$

$d_e(z') > 2C_2 |\tau|^n \max(d_e(x), d_e(y))$. But z'_1 and z'_2 must then also satisfy $|z'_1|, |z'_2| < K + L < |\operatorname{tr} R|^m$. Then there is $w \in \sigma_R^{-1}(\beta^{-1}(z))$ with $l(w) \leq 2^{m+1} + 4m - 1$ by Lemma 33, that is to say $|\beta^{-1}(z)| \leq 2^{m+1} + 4m - 1$. Then, by Proposition 20, $\|\beta^{-1}(z')\| \leq 2^{m+1} + 4m + D - 1$. Letting $r = \lfloor L / (2^{m+1} + 4m + D - 1) \rfloor - 1$, where the brackets represent the greatest integer function, the result now follows from Proposition 6 if we take $f(z') = \|\beta^{-1}(z')\|$. It is straightforward to check that $\lim_{m \rightarrow \infty} r = \infty$, since $|\operatorname{tr} R| > 2$.

The proof for $|\operatorname{tr} R| \leq 2$ is analogous; just replace $|\operatorname{tr} R|$ by 3 or 6 as needed and use the other half of Lemma 33. $\square$

REFERENCES

- [1] O.V. Bogopol'skii. Infinite commensurable hyperbolic groups are bi-Lipschitz equivalent. *Algebra and Logic*, 36(3):155–63, 1997.
- [2] S. Cleary and T.R. Riley. A finitely presented group with unbounded dead-end depth. *Proceedings of the American Mathematical Society*, 134(2):343–9, 2006. [arXiv:math.GR/0406443](#).
- [3] S. Cleary and J. Taback. Dead end words in lamplighter groups and other wreath products. *The Quarterly Journal of Mathematics*, 56(2):165–78, 2005. [arXiv:math.GR/0309344](#).
- [4] D.B.A. Epstein et al. *Word Processing in Groups*. Jones and Bartlett, Boston, 1992.
- [5] T. Riley and A.D. Warshall. The unbounded dead-end depth property is not a group invariant. To appear in *International Journal of Algebra and Computation*, [arXiv:math.GR/0504121](#).
- [6] W. Thurston. *Three-dimensional Geometry and Topology*, Vol. 1 (S. Levy, ed.). Princeton Mathematical Series, vol. 35. Princeton University Press, Princeton, 1997.

YALE UNIVERSITY, DEPARTMENT OF MATHEMATICS, P.O. BOX 208283, NEW HAVEN, CT 06520-8283, USA

E-mail address: andrew.warshall@yale.edu